

CONgenital CORvitia

Project group

B.J.M. Mulder, AMC Amsterdam
 E.T. van der Velde, LUMC Leiden
 J.W.J. Vriend, ICIN
 C.J.M. Engelfriet-Rijk, ICIN
 I. Harms, ICIN
 H.D.M. Murk Jansen, ICIN
 L.M. Roffel, ICIN
 W. Hoekstra, ICIN



National registry and DNA-bank of adults
 with congenital heart disease

www.CONCOR.net



Steering committee

B.J.M. Mulder, chairman, ICIN
 E.T. van der Velde, LUMC Leiden
 E.C.M. Mariman, AZM Maastricht
 I. Kamerling, Netherlands Heart Foundation
 F.J. Meijboom, EMC Rotterdam
 H.W. Vliegen, LUMC Leiden
 G. Veen, VUMC Amsterdam
 P.G. Pieper, AZG Groningen
 H.W.M. Plokker, UMCU Utrecht
 A.P.J. van Dijk, UMC St. Radboud Nijmegen
 O. Daniëls, UMC St. Radboud Nijmegen
 J.L.M. Stappers, AZM Maastricht
 J.G.P. Tijssen, AMC Amsterdam

Number of included patients:
4452

1 November 2004




Figure 1. Number of patients included.




Figure 2. Most frequent main diagnoses.

Table 1. Most frequent main diagnoses and interventions with arrhythmia.

Main diagnosis	Number of patients (prevalence per diagnosis)	
	Supraventricular	Ventricular only
Tetralogy of Fallot	111 (18.2%)	21 (3.4%)
Transposition of the great arteries	108 (33.8%)	5 (1.6%)
ASD	121 (30.5%)	2 (0.5%)
ASD I	36 (22.5%)	0
Aortic coarctation	25 (5.6%)	3 (0.7%)
Tricuspid atresia	26 (49.1%)	0
Functionally univentricular heart	21 (42.0%)	0
Congenitally corrected transposition	20 (27.4%)	5 (6.8%)
Other	Supraventricular	Ventricular only
Mustard/Senning	103 (39.5%)	3 (1.1%)
Fontan	53 (46.9%)	1 (0.9%)




Figure 3. Age distribution (red) and prevalence of arrhythmia (blue) per age group.